

# UNCLASSIFIED

<b>AD NUMBER</b>
ADB206148
<b>NEW LIMITATION CHANGE</b>
<b>TO</b> Approved for public release, distribution unlimited
<b>FROM</b> Distribution authorized to DoD only; Specific Authority; 24 Jan 95. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, Attn; MCMR-RMI-S, Ft. Detrick, MD 21702-5012.
<b>AUTHORITY</b>
U.S. Army Medical Research and Materiel Command ltr., dtd January 21, 2000.

THIS PAGE IS UNCLASSIFIED

AD \_\_\_\_\_

CONTRACT NUMBER: DAMD17-95-C-5046



TITLE: Antibody DNA Vaccine Against Malaria

PRINCIPAL INVESTIGATOR: Maurizio Zanetti, M.D.

CONTRACTING ORGANIZATION: Molecular Medicine Technologies Inc.  
San Diego, California 92101

REPORT DATE: October 1995

TYPE OF REPORT: Final, Phase I

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to DoD  
Components only, Specific Authority. Other requests shall be  
referred to the Commander, U.S. Army Medical Research and  
Materiel Command, ATTN: MCMR-RMI-S, Fort Detrick, MD 21702-5012

The views, opinions and/or findings contained in this report are  
those of the author(s) and should not be construed as an official  
Department of the Army position, policy or decision unless so  
designated by other documentation.

DTIC QUALITY INSPECTED 3

19960111 064

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 1995	3. REPORT TYPE AND DATES COVERED Final, Phase I 15 Mar 95 - 14 Sep 95	
4. TITLE AND SUBTITLE Antibody DNA Vaccine Against Malaria			5. FUNDING NUMBERS DAMD17-95-C-5046	
6. AUTHOR(S) Maurizio Zanetti, M.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Molecular Medicine Technologies Inc. San Diego, California 92101			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Distribution authorized to DoD components only, Specific Authority. Other request shall be referred to the Commander, U.S. Army Medical Research and Materiel Command, ATTN: MCMR-RMI-S, Fort Detrick, MD 21702-5012			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The overall goal of this contract was to study the characteristics of the immune response against a malaria parasite antigen using vaccination with DNA. During the tenure of the contract we have analyzed in great detail all the immunologic aspects we had anticipated we would investigate. Collectively, the findings indicate the following: a) A <u>single</u> intraspleen inoculation of DNA of an antigenized antibody H chain coding for a malaria parasite is sufficient to elicit immunity against the malaria. The initial antibody response is primarily IgM. b) A <u>single</u> DNA injection is also sufficient to induce immunologic memory. Following protein booster injection (challenge) a vigorous secondary antibody response develops with isotype switch to IgG1. c) The initial IgM antibodies can be switched to IgG1 by immunizing with a gene chimerized with the DNA coding for the cytokine GM-CSF. d) DNA immunization induces antibodies that react with <i>P. falciparum</i> sporozoites, hence suggesting that the immunogenic process mimics natural immunity by the parasite. We deem that these results are new and have brought new understanding on how immunity to the malaria parasite may work and how DNA immunization can be exploited to program the immune system to mount an effective immune response.				
14. SUBJECT TERMS DNA Vaccination; Malaria Parasite; Antibodies; NANP			15. NUMBER OF PAGES 15	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT limited	

## GENERAL INSTRUCTIONS FOR COMPLETING SF 298

The Report Documentation Page (RDP) is used in announcing and cataloging reports. It is important that this information be consistent with the rest of the report, particularly the cover and title page. Instructions for filling in each block of the form follow. It is important to **stay within the lines** to meet **optical scanning requirements**.

**Block 1. Agency Use Only (Leave blank).**

**Block 2. Report Date.** Full publication date including day, month, and year, if available (e.g. 1 Jan 88). Must cite at least the year.

**Block 3. Type of Report and Dates Covered.** State whether report is interim, final, etc. If applicable, enter inclusive report dates (e.g. 10 Jun 87 - 30 Jun 88).

**Block 4. Title and Subtitle.** A title is taken from the part of the report that provides the most meaningful and complete information. When a report is prepared in more than one volume, repeat the primary title, add volume number, and include subtitle for the specific volume. On classified documents enter the title classification in parentheses.

**Block 5. Funding Numbers.** To include contract and grant numbers; may include program element number(s), project number(s), task number(s), and work unit number(s). Use the following labels:

<b>C</b> - Contract	<b>PR</b> - Project
<b>G</b> - Grant	<b>TA</b> - Task
<b>PE</b> - Program Element	<b>WU</b> - Work Unit Accession No.

**Block 6. Author(s).** Name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. If editor or compiler, this should follow the name(s).

**Block 7. Performing Organization Name(s) and Address(es).** Self-explanatory.

**Block 8. Performing Organization Report Number.** Enter the unique alphanumeric report number(s) assigned by the organization performing the report.

**Block 9. Sponsoring/Monitoring Agency Name(s) and Address(es).** Self-explanatory.

**Block 10. Sponsoring/Monitoring Agency Report Number.** (If known)

**Block 11. Supplementary Notes.** Enter information not included elsewhere such as: Prepared in cooperation with...; Trans. of...; To be published in.... When a report is revised, include a statement whether the new report supersedes or supplements the older report.

**Block 12a. Distribution/Availability Statement.** Denotes public availability or limitations. Cite any availability to the public. Enter additional limitations or special markings in all capitals (e.g. NOFORN, REL, ITAR).

**DOD** - See DoDD 5230.24, "Distribution Statements on Technical Documents."

**DOE** - See authorities.

**NASA** - See Handbook NHB 2200.2.

**NTIS** - Leave blank.

**Block 12b. Distribution Code.**

**DOD** - Leave blank.

**DOE** - Enter DOE distribution categories from the Standard Distribution for Unclassified Scientific and Technical Reports.

**NASA** - Leave blank.

**NTIS** - Leave blank.

**Block 13. Abstract.** Include a brief (*Maximum 200 words*) factual summary of the most significant information contained in the report.

**Block 14. Subject Terms.** Keywords or phrases identifying major subjects in the report.

**Block 15. Number of Pages.** Enter the total number of pages.

**Block 16. Price Code.** Enter appropriate price code (*NTIS only*).

**Blocks 17. - 19. Security Classifications.** Self-explanatory. Enter U.S. Security Classification in accordance with U.S. Security Regulations (i.e., UNCLASSIFIED). If form contains classified information, stamp classification on the top and bottom of the page.

**Block 20. Limitation of Abstract.** This block must be completed to assign a limitation to the abstract. Enter either UL (unlimited) or SAR (same as report). An entry in this block is necessary if the abstract is to be limited. If blank, the abstract is assumed to be unlimited.

## FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

N/A Where copyrighted material is quoted, permission has been obtained to use such material.

N/A Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

✓ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

✓ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

N/A For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

N/A In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

N/A In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories. ...

Accession For ...	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
E-4	

  
PI - Signature      10/13/1995  
Date

## (5) INTRODUCTION

Malaria continues to be a major threat to man kind in many areas of the globe, particularly in underdeveloped countries. *Plasmodium falciparum*, one of the four etiologic agents of malaria in humans, is widely distributed and causes severe, and frequently fatal, malaria in children and nonimmune adults. Its control is hampered by the increased resistance to chemoprophylaxis, chemotherapy as well as eradication of the Anopheles mosquito vectors with insecticides.

The major surface proteins of malaria sporozoites, the circumsporozoites (CS) proteins, have unusual structural and immunological properties. It has now been established that in all *plasmodium* species the central portion of the molecule consists of a repetitive amino acid sequence which contains the protein immunodominant B-cell epitope(s). In *P. falciparum* the central domain contains about 40 repeat of the tetrapeptide Asn-Ala-Asn-Pro (NANP) plus 3 to 4 Asn-Val-Asp-Pro (NVDP) repeats. The presence of internal repeats likely derives from tandem duplication and is considered to have structure-function relevance.

To date there are no efficient methods to immunize against this parasite and much of what is done towards the development of a malaria vaccine is still experimental in nature. In particular, in addition to identify new parasite antigens that may be suitable for immunization and protection, emphasis is also placed on new methods to vaccinate and present antigen to the immunocompetent host.

The goal of our original proposal was to validate a new principle and method of immunization against the malaria parasite, gene immunization, using as a vaccine an immunoglobulin gene re-engineered to encode a discrete epitope of the malaria parasite, "antigenized antibody gene". As a model system we will use the tetrapeptide sequence NANP of the malaria parasite *Plasmodium falciparum* CS protein.

The studies proposed made use plasmid DNA ("naked" DNA) for the heavy chain of an antibody antigenized to express (NANP)<sub>3</sub>. This gene was used in pilot studies to immunize mice with the intent of inducing anti-NANP antibodies. The studies were designed at verifying feasibility by empirically establishing a series of parameters such as specificity and titer of the antibody response, isotype of NANP-reactive antibodies, duration of the antibody response and priming effect on a subsequent immune response elicited by a protein antigen. The results obtained in the course of these pilot studies (six months) will be reviewed hereunder.

## (6) BODY

The work was done using a series of plasmid vectors designed and manufactured in the laboratory. These were used to inoculate C57Bl/6 mice to study the immune response that was elicited. Therefore, prior to summarizing the data we will briefly describe the material and methods used throughout the studies.

### Material and Methods:

The plasmid DNA

- (1) γ1NANP. This is a modified version of the γ1WT plasmid (*infra*) where the third hypervariable region was modified to encode three repeats of the tetrapeptide Asn-Ala-Asn-Pro (NANP) sequence. This vector also encodes a genomic human γ1 gene human constant region gene and contains a neomycin resistance gene under the control of the SV40 promoter for the selection of stable transformant cells.

- (2)  $\gamma$ 1WT. This defines plasmid pN $\gamma$ 162 which contains the 2.3 Kb EcoRI DNA fragment carrying the genomic rearrangement of the murine V $H$ <sup>62</sup> gene together with a genomic human  $\gamma$ 1 gene human constant region gene.
- (3) pSV2neo. This is the original plasmid forming the backbone of the pN $\gamma$ 1 vector without the genomic human  $\gamma$ 1 gene human constant region gene.

All plasmid DNAs used for injection were prepared from DH5 $\alpha$  *Escherichia coli* according to standard procedures. The DNA is stored at -20°C until use.

#### Serological assays

- Detection of serum antibodies. This was done on sera collected at regular day intervals. The detection of antibodies to  $\gamma$ 1NANP was done on a 96-well polyvinyl microtiter plates (Dynatech, Gentilly, VA) coated (2.5  $\mu$ g/ml) by drying at 37°C. After coating, wells were blocked with 1% bovine serum albumin (BSA) in phosphate buffered saline (PBS) 0.15 M, pH 7.3. Wells were incubated overnight at 4°C with mice sera diluted in 1% BSA-PBS, containing 1% Tween-20 (PBSA). The bound antibodies were revealed using a goat antibody to mouse  $\gamma$ -globulin absorbed with human  $\gamma$ -globulins and conjugated with horseradish peroxidase (HP) (Sigma, St. Louis, MO). The bound peroxidase was revealed by adding o-phenylenediamine dihydrochloride and H<sub>2</sub>O<sub>2</sub>. Plates were read after 30 minutes in a micro-plate reader (Vmax, Molecular Devices) at 492 nm. Tests were done in duplicate. The detection of antibodies against the NANP peptide was done similarly using microtiter wells coated with the synthetic peptide KK(NANP)<sub>3</sub> (2.5  $\mu$ g/ml). As a control, we used a peptide of human vitronectin. In the experiments reported herein, the peptide was diluted in 0.9% NaCl and adsorbed on the plate by drying at 37°C. (Experiments that will be reported in the future will be done using a more gentle approach of binding the peptide to the plate, that is a solution in carbonate buffer 0.01 M at pH 9.6).
- Detection of the Transgene Product. This was done on sera done on sera collected at regular day intervals. The presence of  $\gamma$ 1NANP H chain transgene polypeptide in the serum of mice as detected using a capture assay. Briefly, 1:10 dilution of individual mouse sera in PBSA were incubated on 96-well plate coated with a goat antibody to human  $\gamma$ -globulin (10  $\mu$ g/ml). To determine the concentration of the transgene product in the serum, O.D. values were plotted against a standard curve constructed with known amounts of human  $\gamma$ -globulins diluted in PBSA containing 10% normal mouse serum. The bound antibodies were revealed using a HP-conjugated goat antibody to human  $\gamma$ -globulin (H chain specific) absorbed with murine immunoglobulins (Sigma, St. Louis, MO). The bound peroxidase activity was revealed by adding o-phenylenediamine dihydrochloride and H<sub>2</sub>O<sub>2</sub>. Because antibody titers were found only in the groups immunized intraspleen, the transgene was searched for only in these groups. Tests were done in duplicate.
- Isotype analysis of the immune response. Ig class determination was done by ELISA using isotype-specific rabbit antibodies (Mouse Typer Subtyping Kit, Bio-Rad, Hercules, CA) used at the dilution recommended by the manufacturer.

#### Molecular biology techniques

The expression of the transgenic mRNA was studied in the spleen of an immunized mouse inoculated via the intrasplenic route by reverse PCR. Briefly, the spleen was harvested on day 20 after the initial inoculation, frozen at -170 °C. The cells were prepared by gentle teasing. Typically, PolyA<sup>+</sup> mRNA was isolated from 11 mg of spleen

tissue using the the Micro-FastTrack mRNA isolation kit (Invitrogen, San Diego, CA). cDNA was synthesized by reverse transcription using the cDNA Cycle Kit (Invitrogen, San Diego). Specific PCR primers for the murine  $V_H$  gene and the human  $\gamma 1$  C region gene (the transgenes) were designed using Oligo 4.0 Primer Analysis Software (NBI, Inc., Plymouth, MN) from the sequences of the murine  $V_H^{62}$  and human  $\gamma 1$  genes obtained from the National Center for Biotechnology Information (NCBI) GenBank (Bethesda, MA). The PCR reaction conditions were as follows: 1 min. at 92 °C followed by 2 min. at 40 °C and 3 min. at 72 °C. This cycle was repeated 30 times. cDNA amplification and size determination (  $V_H$  = 240 bp; human  $\gamma 1$  C region gene = 466 bp) were checked by electrophoresis using 4% NuSieve 3:1 agarose gel (FMC, Rockland, ME). To generate sticky ends on the PCR products we used the restriction half-site strategy where the oligonucleotide primers used contained three nucleotides (TTC) of the 6-bases recognition site of EcoRI site at their 5' end. The fragments were then cloned in pBluescript KS vector (Stratagene, San Diego, CA) for sequencing by the dideoxy method Sequenase 2.0 DNA sequencing Kit (USB, Cleveland OH). For Southern blot studies the cDNA was analyzed after electrophoresis on a 1% agarose gel and transfer to Hybond-N<sup>+</sup> nylon membrane (Amersham). The Blot was hybridized with oligonucleotide probes labeled using T4 polynucleotide kinase forward reaction in presence of  $\gamma$ ATP<sup>32</sup>.

#### Histology of spleens after a single DNA inoculation.

Spleens were fixed in 4% paraformaldehyde for 24 hours, rinsed with PBS and included in Tissue-Tek (Miles, Elkhart, IN) and frozen at -70 °C. Blocks of frozen tissue were cut (5  $\mu$ m) on a cryostat and stained with hematoxiniln and eosin. Sections were inspected by two investigators.

### Results

The results collected over the past six months can be summarized as follows:

#### 1) Effect of the route of DNA inoculation

Only the intra-spleen route of immunization yields antibodies to both the  $\gamma 1$ NANP protein and the synthetic peptide NANP. The response was specific in that none of the mice immunized with control DNA had a measurable antibody response. Notably, control mice primed by inoculation intra-spleen of antigenized antibody  $\gamma 1$ NANP mounted a weak primary antibody response. These results are summarized in Table 1.

**Table 1. The Route of Inoculation Determines the fate of DNA Immunization**

Route of Inoculation	No. of Mice	No. of Injections	Antibody Titers (Log)		
			Day 0	Day 14	Day 35
Intramuscular	5	4	$\leq 2.3$	2.4 $\pm$ 0.2	2.4 $\pm$ 0.2
Subcutaneous	5	4	$\leq 2.3$	$\leq 2.3$	$\leq 2.3$
Intravenous	5	4	$\leq 2.3$	$\leq 2.3$	$\leq 2.3$
Intradermal	4	3	$\leq 2.3$	$\leq 2.3$	2.4 $\pm$ 0.2
Intra-spleen	5	1	$\leq 2.3$	<b>3.1<math>\pm</math>0.4</b>	<b>3.1<math>\pm</math>0.2</b>
Intra-spleen + Intravenous	5	1 + 3	$\leq 2.3$	2.8 $\pm$ 0.2	2.8 $\pm$ 0.2



2) Production of transgene immunoglobulins

Transgene immunoglobulins were detected in the serum of all mice immunized intraspleen. These results are summarized below in Table 2.

**Table 2. Detection of transgene immunoglobulins in the serum of immunized mice**

Route	Antigen	No. of Mice	No. of Injections	Transgene Product (ng/ml)		
				Day 0	Day 7	Day 14
i.s.	$\gamma$ 1NANP (DNA)	10	1	Undetectable	3.5 $\pm$ 3.2	11.8 $\pm$ 5.6
i.s.	pSV2Neo (DNA)	4	1	Undetectable	Undetectable	Undetectable
i.s.	$\gamma$ 1NANP (protein)	4	1	Undetectable	6 $\pm$ 5	Undetectable
i.s.	OVA	4	1	Undetectable	Undetectable	Undetectable

3) The antibody response

The response anti-transgene immunoglobulins and anti-NANP peptide is specific for DNA immunization into the spleen. These data are summarized in Tables 3 and 4. Notably, mice inoculated intraspleen with 50  $\mu$ g of equivalent transgene immunoglobulin did not make any measurable response.

**Table 3. Detection of Anti- $\gamma$ 1NANP Antibodies in Mice Immunized Intraspleen**

Immunogen	No. of Mice	No. of Injections	No. of Responders	Antibody Titers (Log)			
				day 0	14	28	53
$\gamma$ 1NANP (DNA)	8	1	8/8	2.3	2.8	2.9	2.9
pSV2Neo (DNA)	4	1	0/4	2.3	2.3	2.3	2.4
$\gamma$ 1NANP (protein)	4	1	3/4	2.3	2.4	2.4	2.5
OVA (protein)	4	1	0/4	2.3	2.3	2.3	2.3

**Table 4. Detection of Anti-Peptide Antibodies in Mice Immunized Intraspleen**

Immunogen	No. of Mice	No. of Injections	No. of Responders	Antibody Titers (Log)			
				day 0	14	28	53
$\gamma$ 1NANP (DNA)	8	1	8/8	2.3	2.5	2.8	2.8
pSV2Neo (DNA)	4	1	0/4	2.3	2.3	2.3	2.3
$\gamma$ 1NANP (protein)	4	1	0/4	2.3	2.3	2.3	2.3
OVA (protein)	4	1	0/4	2.3	2.3	2.3	2.3

4) Induction of immunologic memory

Intraspleen DNA immunization induces immunologic memory. We verified that mice primed by virtue of a single intraspleen DNA inoculation had acquired immunologic memory as defined by the ability to mount a vigorous response upon challenge with a protein antigen. These data are summarized in Table 5 and 6.

**Table 5. Detection of antibodies to  $\gamma$ 1NANP in mice primed intra-spleen and boosted subcutaneously**

Priming	Booster	No. of Mice	Day 0	Day 14	Day 28
$\gamma$ 1NANP DNA	$\gamma$ 1NANP DNA	4	2.8 $\pm$ 0.4	2.9 $\pm$ 0.1	2.9 $\pm$ 0.1
$\gamma$ 1NANP DNA	$\gamma$ 1NANP protein	4	2.5 $\pm$ 0.2	3.4 $\pm$ 0.2	3.5 $\pm$ 0.3
pSVneo DNA	$\gamma$ 1NANP protein	4	2.3 $\pm$ 0	2.4 $\pm$ 0.2	2.5 $\pm$ 0.2
$\gamma$ 1NANP protein	$\gamma$ 1NANP protein	4	2.5 $\pm$ 0.2	2.9 $\pm$ 0.2	2.9 $\pm$ 0.2
OVA protein	OVA protein	4	2.3 $\pm$ 0	2.3 $\pm$ 0	2.3 $\pm$ 0

Titers are expressed as means  $\pm$  SD of Log values of individual mice. Day 0 refers to the day of the booster immunization, that is 200 days after the initial immunization. DNA booster (100  $\mu$ g) was given in saline solution intravenously. Protein booster (50  $\mu$ g) was given in IFA subcutaneously.


**Table 6. Detection of antibodies to synthetic peptide NANP in mice primed intra-spleen and boosted subcutaneously**

Priming	Booster	No. of Mice	Day 0	Day 14	Day 28
$\gamma$ 1NANP DNA	$\gamma$ 1NANP DNA	4	2.9 $\pm$ 0	2.9 $\pm$ 0	2.9 $\pm$ 0
$\gamma$ 1NANP DNA	$\gamma$ 1NANP protein	4	3 $\pm$ 0.2	3.6 $\pm$ 0.3	3.7 $\pm$ 0.4
pSVneo DNA	$\gamma$ 1NANP protein	4	2.3 $\pm$ 0	2.3 $\pm$ 0	2.3 $\pm$ 0
$\gamma$ 1NANP protein	$\gamma$ 1NANP protein	4	2.4 $\pm$ 0.3	2.5 $\pm$ 0.4	2.6 $\pm$ 0.6
OVA protein	OVA protein	4	2.3 $\pm$ 0	2.3 $\pm$ 0	2.3 $\pm$ 0

Titers are expressed as means  $\pm$  SD of Log values of individual mice. Day 0 refers to the day of the booster immunization, that is 200 days after the initial immunization. DNA booster (100  $\mu$ g) was given in saline solution intravenously. Protein booster (50  $\mu$ g) was given in IFA subcutaneously. A synthetic peptide from the sequence of human vitronectin was used as a control. The average reactivity on control peptide was reproducibly 2.3.

As indicated a single booster injection with the  $\gamma$ 1NANP protein in mice primed with DNA intra-spleen augmented markedly the antibody response to both the transgene product as well as against the NANP peptide. The effect was specific in that none of the other groups (DNA  $\rightarrow$  DNA; or protein intra-spleen  $\rightarrow$  protein; or control DNA  $\rightarrow$  protein) had such a marked antibody response. It is worth noting that a booster injection with DNA  $\gamma$ 1NANP failed to induce an increase in the antibody titer over the primary immune response.


The antibody response (time course and relative binding) is also shown in Figure 1.



**Figure 1.** Time course and effect of booster immunization on the immune response to  $\gamma$ 1NANP (top) and NANP peptide (bottom). Sera were tested individually at 1:1600 dilution. Experimental groups are as described in the text. Time points (week) on left refer to bleeds after the primary injection. Time points on the right correspond to bleeds after the booster immunization. Time 0 refers to the day of the booster injection.

##### 5) Isotype analysis of the immune response during the primary and secondary response

Much to our surprise we found that the antibodies produced in response to a single intra-spleen immunization are prevalently of the IgM isotype. This suggested that priming of the immune response by intra-spleen DNA immunization may occur in conditions of limited production of cytokines, including those necessary for isotype switching. We also found that after booster, mice immunized once intraspleen with DNA undergo rapid IgM->IgG1 switch as shown below in Figure 2.



**Figure 2.** Isotype analysis of the immune response in mice primed intra-spleen with  $\gamma$ 1NANP DNA (Left) or  $\gamma$ 1NANP protein (Right), and boosted subcutaneously with the transgene protein in IFA. Symbols are indicated in the Figure. Sera were tested as a pool at a 1:1600 dilution. The first two columns in each group refer to day 0 and week 4 after priming; the last two columns refer to the day of the booster injection (Day 0) and week 4 after the booster, respectively.

The results indicate that after booster with the transgene protein mice make a strong antibody response of the IgG1 and to a lesser extent IgG2b isotype, but no significant increase of the IgM response over the primary immunization is observed. In contrast, mice primed intra-spleen with the transgene protein ( $\gamma$ 1NANP) make a primary IgG1 response but not IgG2b antibodies.

Therefore, a booster immunization with protein in animals primed with DNA via the intra-spleen route readily induced isotype switch and production of IgG1 antibodies much greater than in the group primed with the transgene protein directly into the spleen. Intra-spleen immunization with DNA induces IgG2b antibodies in the secondary response. Because these effects were of more limited magnitude in mice intra-spleen with the antigenized antibody, we conclude that these effects are not solely the result of the booster immunization but also of a state of immunologic memory established by priming with DNA intra-spleen.

6. Strategies of DNA immunization intraspleen to modulate the quality of the immune response

Isotype analysis of the primary and secondary antibody responses indicates that IgM is the prevalent isotype during the primary immune response following a single intra-spleen inoculation of DNA. This has suggested to us that the T cell help provided is insufficient to promote switch. Studies of the secondary immune response, however, indicate that the initial IgM response switches to IgG1 upon new contact with antigen.

For the purpose of vaccination it may be desirable to induce IgG1 antibodies at the outset of immunization. In immunization with protein antigens this is achieved by incorporating the antigen in an immunologic adjuvant (e.g., CFA, alum etc.). Induction of an IgG1 response by immunization with naked DNA may require multiple DNA injections.

There are, however, possibilities to direct isotype switch at the time of immunization without using conventional immunologic adjuvants. For instance, the use of cytokines at the time of immunization or the use of antigen/cytokine chimeras may be of advantage to immunization. Relevant to our experiment is the observation by R. Levy and colleagues (Stanford University) that an antibody/GM-CSF chimera induces a marked primary IgG1 anti-idiotype response.

Although these new studies were not contemplated in the original SBIR proposal, we decided to use as the immunogen DNA (e.g.,  $\gamma$ 1NANP) chimerized with GM-CSF or IL-2. We studied the effect(s), if any, of DNA chimeras on: *i*) the isotype response, and *ii*) the quantitative aspects of the overall response. The results obtained are shown in Table 7, 8 and 9.

**Table 7. Induction of IgG1 antibodies during the primary response after intraspleen inoculation with  $\gamma$ 1NANP/GM-CSF DNA chimera**

Exp. #	Priming (DNA)	Booster	No. of Mice	Primary Response (Days)		
				0	14	28
I	$\gamma$ 1NANP	$\gamma$ 1NANP	6	0.239	0.308	0.315
	$\gamma$ 1NANP/GM-CSF	$\gamma$ 1NANP	6	0.217	0.967	0.819
	$\gamma$ 1NANP/IL-2	$\gamma$ 1NANP	6	0.237	0.241	0.233
II	$\gamma$ 1NANP	$\gamma$ 1NANP	3	0.240	0.278	0.320
	$\gamma$ 1NANP/GM-CSF	$\gamma$ 1NANP	3	0.273	0.847	0.638
	$\gamma$ 1NANP/IL-2	$\gamma$ 1NANP	4	0.218	0.283	0.222
III	$\gamma$ 1NANP	$\gamma$ 1NANP	4	0.250	0.283	0.305
	$\gamma$ 1NANP/GM-CSF	$\gamma$ 1NANP	4	0.240	0.791	0.617
	$\gamma$ 1NANP/IL-2	$\gamma$ 1NANP	4	0.282	0.283	0.279

Values correspond to ELISA readings (A492) of pool of sera collected on the day indicated in the Table and tested at 1:1600 dilution.

**Table 8. The antibody response against  $\gamma$ 1NANP protein in mice inoculated intra-spleen with  $\gamma$ 1NANP/cytokine chimera is similar to that elicited by  $\gamma$ 1NANP alone**

DNA Immunogen	Day 0	Day 14	Day 35	Day 14 After boost	Day 28 After boost
$\gamma$ 1NANP/GM-CSF	2.6	3.2 $\pm$ 0.3	3.2 $\pm$ 0.2	3.7 $\pm$ 0.3	4 $\pm$ 0.6
$\gamma$ 1NANP/IL-2	2.6	2.8 $\pm$ 0.2	2.8 $\pm$ 0.4	3.6 $\pm$ 0.2	3.6 $\pm$ 0.2
$\gamma$ 1NANP	2.6	3.2 $\pm$ 0.1	3.2 $\pm$ 0.3	3.8 $\pm$ 0.2	4.1 $\pm$ 0.6

Mice were immunized by a single inoculation of DNA intra-spleen (100  $\mu$ g). A booster injection was given on day 40 by injection of  $\gamma$ 1NANP protein (50  $\mu$ g) in IFA subcutaneously. Values represent the means  $\pm$  SD of 4 mice/group tested individually.

**Table 9. Antibody NANP peptide in mice inoculated intra-spleen with  $\gamma$ 1NANP/cytokine chimera**

DNA Immunogen	Day 0	Day 14	Day 35	Day 14 After boost	Day 28 After boost
$\gamma$ 1NANP/GM-CSF	2.3	2.6 $\pm$ 0.2	2.8 $\pm$ 0.2	3 $\pm$ 0.2	3 $\pm$ 0.1
$\gamma$ 1NANP/IL-2	2.3	2.5 $\pm$ 0.2	2.6 $\pm$ 0.2	2.6 $\pm$ 0.1	2.6 $\pm$ 0.1
$\gamma$ 1NANP	2.3	2.5 $\pm$ 0.2	2.6 $\pm$ 0.2	2.7 $\pm$ 0.2	2.7 $\pm$ 0.2
$\gamma$ 1NANP/GM-CSF	2.3	2.7 $\pm$ 0.2	2.8 $\pm$ 0.2	3.1 $\pm$ 0.2	3.1 $\pm$ 0.21
$\gamma$ 1NANP/IL-2	2.3	2.7 $\pm$ 0.3	2.7 $\pm$ 0.3	3.1 $\pm$ 0.3	3.1 $\pm$ 0.3
$\gamma$ 1NANP	2.3	2.6	2.6	2.9	2.9
$\gamma$ 1NANP/GM-CSF	2.3	2.8 $\pm$ 0.2	2.8 $\pm$ 0.2	3 $\pm$ 0.3	3.1 $\pm$ 0.3
$\gamma$ 1NANP/IL-2	2.3	2.8 $\pm$ 0.3	2.8 $\pm$ 0.3	3 $\pm$ 0.2	3 $\pm$ 0.2
$\gamma$ 1NANP	2.3	2.7 $\pm$ 0.3	2.8 $\pm$ 0.2	3.1 $\pm$ 0.2	3 $\pm$ 0.2

As it can be seen GM-CSF does not appear to have an effect on the quantitative aspect of the anti-peptide response either before or after booster. We conclude that while the addition of GM-CSF produced a marked and reproducible effect on the isotype, it was unable to modify the amounts of antibodies produced.

#### 7) Detection of transgene product in mice immunized with $\gamma$ 1NANP/cytokine chimera

Mice immunized with the  $\gamma$ 1NANP/GM-CSF chimera produce IgG1 antibodies from the onset during the primary response. We compared, therefore, the levels of transgene immunoglobulins in the serum of these mice in order to reveal any difference that may help understanding the qualitative effect of GM-CSF. Table 10 summarizes the results.

**Table 10. Transgene immunoglobulins in mice immunized with DNA/cytokine chimera**

IMMUNOGEN	MICE (No.)	Producers	Transgene Product (ng/ml)
$\gamma$ 1NANP	11	10/11	11.5 $\pm$ 8.4
$\gamma$ 1NANP/GM-CSF	13	1/13	2.2
$\gamma$ 1NANP/IL-2	14	3/14	1.2 $\pm$ 0.1

As indicated mice inoculated with the DNA/cytokine chimera did not produce detectable amounts of the transgene. Therefore, the effect on isotype IgM->IgG1 switch can not be attributed to higher amounts of transgene produced.

#### 8) Histology of spleens after a single DNA inoculation

To check for histological abnormalities subsequent to intraspleen immunization, spleens from DNA-inoculated, saline-inoculated and naive mice were collected 3, 7, 14 and 21 days from the time of injection. We did not observe any gross abnormality in the architecture of the organ or infiltrates of inflammatory cells as compared with spleens from naive mice of the same age or from mice given a sham inoculation of sterile saline solution. Immunohistochemical analysis of frozen sections using monoclonal antibodies to B and activated T lymphocytes showed no changes from controls. Collectively, the intraspleen inoculation of DNA did not cause apparent tissue damage nor appreciable changes in the number and tissue distribution of B and T lymphocytes.

#### 9) Detection of anti-sporozoite antibodies in the serum of mice immunized with DNA

In collaboration with the laboratory of Dr. S. Hoffman (Navy Research Laboratory) we have ascertained that antibodies reactive with the NANP peptide by ELISA also react with air dried sporozoites in indirect immunofluorescence.

The results of such studies have been received 24 hours prior to preparing this final report and they will be reviewed only:

- The pool of sera from mice given a single DNA immunization and subsequently boosted with the NANP-containing protein, was positive (Titer 1:40-1:80). The serum of a mouse immunized only once with  $\gamma$ 1NANP DNA was positive (titer 1:20). This suggests that one DNA immunization alone is sufficient to elicit anti-sporozoite antibodies.
- The serum of mice primed with  $\gamma$ 1NANP/GM-CSF or /IL-2 DNA chimeras, and boosted with the NANP-containing protein failed to react by IFA. The discrepancy could be explained on the basis of the isotype produced in response to the different immunization regimens. In fact, priming with the cytokine chimera predisposes to a great diminution of the IgG2b secondary response. Therefore, it is possible that the difference observed may reflect a more profound qualitative change in the antibody response. It should be mentioned that many reports in the mouse have concluded that antibody-mediated protection is mediated by IgG2a antibodies.
- The positive (+) control and the various negative (-) controls they all performed as expected. Therefore, the test was internally consistent.

10) Detection of the transgene mRNA in the spleen of inoculated mice

The cDNA was analyzed by Southern blot after electrophoresis on a 1% agarose gel and transfer to Hybond-N<sup>+</sup> nylon membrane (Amersham). The Blot was hybridized with oligonucleotide probes labeled using T4 polynucleotide kinase forward reaction in presence of  $\gamma$ ATP<sup>32</sup>. A strong hybridization band was seen with the amplification product of this reaction (240 bp) but not with the amplification product of a spleen from an unmanipulated mouse germline gene. This result was confirmed by sequencing the amplified and reverse-transcribed V<sub>H</sub>. This suggests that after intraspleen inoculation the transgene persist in the spleen (B cells ) for a prolonged period of time.

**(7) CONCLUSIONS**

In conclusion, we have answered all the questions posed in the contract under whose tenure the present work was performed.

1. The main point of our studies is that a single intraspleen inoculation of DNA of an antigenized antibody H chain coding for a malaria parasite is sufficient to elicit immunity against the malaria. The initial antibody response is primarily IgM.
2. A single DNA injection is also sufficient to induce immunologic memory. As shown following a protein booster injection (challenge), a vigorous secondary antibody response develops with isotype switch to IgG1.
3. The initial IgM can be switched to IgG1 by immunizing with a gene chimerized with the DNA coding for GM-CSF.
4. DNA immunization induces antibodies that react with *P. falciparum* sporozoites, hence suggesting that the immunogenic process mimics natural immunity by the parasite.

Future work will have to better define the immunological parameters to maximize the immunogenicity with one injection only as well as to identify the methods to produce high levels protective antibodies, e.g., correct isotype.

**(8) REFERENCES**

There is very little published so far on the induction of immunity against malaria using DNA vaccination. The one relevant publication is:

Sedegah, M., Hedstrom, R., Hobart, P. and Hoffman, S. L. 1994. Protection against malaria by immunization with plasmid DNA encoding circumsporozoite protein. *Proc. Natl. Acad. Sci. USA* 91:9866.

Additional references related to current approaches to DNA vaccination dealing with the induction of an antibody response are listed below. It should be noted that in most if not all instances induction of antibodies required multiple injections of DNA. Our study show that one DNA injection is sufficient to elicit an antibody response directed against the malaria peptide and the malaria parasite (native antigen).

Cox, G., Zamb, T. and Babiuk, L. 1993. Bovine herpesvirus1: Immune responses in mice and cattle injected with plasmid DNA. *J. Virol.* 67:5664.



Davis, H. L., Michel, M. L., Mancini, M., Schleef, M. and Whalen, R. G. 1994. Direct gene transfer in skeletal muscle: plasmid DNA-based immunization against the hepatitis B virus surface antigen. *Vaccine* 12:1503.

Fynan, E. F., Webster, R. G., Fuller, D. H., Haynes, J. R., Santoro, J. C. and Robinson, H. L. 1993. DNA vaccines: protective immunizations by parenteral, mucosal and gene-gun inoculations. *Proc. Natl. Acad. Sci. USA* 90:11478.

Raz, E., Carson, D. A., Parker, S. E., Parr, T. B., Abai, A. M., Aichinger, G., Gromkowski, S. H., Singh, M., Lew, D., Yankauckas, M. A., Baird, S. M. and Rhodes, G. H. 1994. Intradermal gene immunization: the possible role of DNA uptake in the induction of cellular immunity to viruses. *Proc. Natl. Acad. Sci. USA* 91:9519.

Robinson, H., Hunt, L. and Webster, R. 1993. Protection against a lethal influenza virus challenge by immunization with a haemagglutinin-expressing plasmid. *Vaccine* 11:957.

Ulmer, J. B., Donnelly, J. J., Parker, S. E., Rhodes, G. H., Felgner, P. L., Dwarki, V. J., Gromkowski, S. H., Deck, R. R., Dewitt, C. M., Friedman, A., Hawe, L. A., Leander, K. R., Martinez, D., Perry, H. C., Shiver, J. W., Montgomery, D. L. and Liu, M. A. 1993. Heterologous protection against influenza by injection of DNA encoding a viral protein. *Science* 259:1745.

Wang, B., Ugen, K., Srikantan, V., Agadjanyan, M., Dang, K., Refaeli, Y., Sato, A., Boyer, J., Williams, W. and Weiner, D. 1993. Gene inoculation generates immune responses against human immunodeficiency virus type 1. *Proc. Natl. Acad. Sci. USA* 90:4156.

Received 2/8/00



DEPARTMENT OF THE ARMY  
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND  
504 SCOTT STREET  
FORT DETRICK, MARYLAND 21702-5012

REPLY TO  
ATTENTION OF:

MCMR-RMI-S (70-1y)

21 Jan 00

MEMORANDUM FOR Administrator, Defense Technical Information  
Center, ATTN: DTIC-OCA, 8725 John J. Kingman  
Road, Fort Belvoir, VA 22060-6218

SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports written for the attached Awards. Request the limited distribution statements for Accession Document Numbers listed be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Virginia Miller at DSN 343-7327 or by email at virginia.miller@det.amedd.army.mil.

FOR THE COMMANDER:

Encl  
as

A handwritten signature in cursive script, reading "Phylis Rinehart", is written over the typed name and title.  
PHYLIS M. RINEHART  
Deputy Chief of Staff for  
Information Management